

# $\alpha$ -Dedekind complete archimedean vector lattices versus $\alpha$ -quasi- $F$ spaces

Anthony J. Macula

Westfield State College, Westfield, MA 01086, USA

Received 17 October 1989

Revised 4 January 1991

## Abstract

Macula, A.J.,  $\alpha$ -Dedekind complete archimedean vector lattices versus  $\alpha$ -quasi- $F$  spaces, *Topology and its Applications* 44 (1992) 217–234.

$\alpha$  denotes an uncountable cardinal number, and  $\mathcal{W}(\alpha)$  denotes the category of archimedean vector lattices with distinguished weak unit and unit preserving vector lattice homomorphisms. In this paper we show that in  $\mathcal{W}(\alpha)$ , the full subcategory of  $\alpha$ -Dedekind complete objects is epireflective. We explain how the Yosida functor connects the algebraic notions of  $\alpha$ -Dedekind complete,  $\alpha$ -dense, and  $\alpha$ -jam-dense with the topological notions of  $\alpha$ -quasi- $F$ ,  $\alpha$ -irreducible, and  $\alpha$ -SpFi morphism. We then go on to show that the  $\alpha$ -quasi- $F$  cover of a compact space  $X$ , denoted  $(QF_\alpha X, q_\alpha)$ , is the Yosida space of the  $\alpha$ -Dedekind complete epireflection of  $C(X)$  in  $\mathcal{W}(\alpha)$ . Finally we show that in the topological category  $\alpha$ -SpFi, the full subcategory of  $\alpha$ -quasi- $F$  spaces is monoreflective, and for each compact space  $X$ ,  $(QF_\alpha X, q_\alpha)$  is the  $\alpha$ -quasi- $F$  monoreflection of  $X$ .

**Keywords:**  $\alpha$ -Dedekind complete vector lattice,  $\alpha$ -quasi- $F$  spaces, Yosida space, spaces with filters,  $\alpha$ -jam-dense.

**AMS (MOS) Subj. Class.:** Primary 06B15, 06A23, 18B30, 54G05, 54C10; secondary 06F15, 18A20.

## 1. Introduction

$\alpha$  denotes an uncountable cardinal number or the symbol  $\infty$ . The meaning of  $\alpha = \infty$  will be clear from the context. We write  $\alpha < \infty$  to mean that  $\alpha$  is an arbitrary cardinal number.

$\mathcal{W}$  denotes the category of archimedean vector lattices with distinguished weak unit and unit preserving vector lattice homomorphisms. An element  $u \in L \in |\mathcal{W}|$  is called a *weak unit* if the band (complete ideal) generated by  $u$  is all of  $L$  [32, 10].  $u$  is called a *strong unit* if the principal ideal generated by  $u$  is all of  $L$ .  $\mathcal{S}$  denotes

the category of archimedean vector lattices with distinguished strong unit and unit preserving vector lattice homomorphisms. Obviously, a strong unit is a weak unit, and  $\mathcal{S}$  is a subcategory of  $\mathcal{W}$ .  $L^+$  denotes all the *strictly* positive elements of  $L$ , i.e.,  $L^+ = \{a \in L: a > 0\}$ .

An archetypical  $\mathcal{W}$  object is  $C(X)$  (the ring of continuous real-valued functions on a topological space  $X$  [11]) with the constant function  $\mathbf{1}$  taken as the distinguished weak unit. Note that the weak unit  $\mathbf{1}$  is indeed a strong unit, and  $(C(X), \mathbf{1})$  is an  $\mathcal{S}$  object.

**Definition 1.1.** A  $\mathcal{W}$  morphism  $\varphi: L \rightarrow M$  is called  $\alpha$ -complete if, for  $A \subset L$  with  $|A| < \alpha$ , we have that  $\varphi(\bigvee^L A) = \bigvee^M \varphi[A]$  whenever  $\bigvee^L A$  exists in  $L$ .  $\mathcal{W}(\alpha)$  denotes the category of  $\mathcal{W}$  objects with  $\alpha$ -complete morphisms.

**Definition 1.2.** An  $L \in |\mathcal{W}|$  is called  $\alpha$ -Dedekind complete (also  $\alpha$ -jamd-complete [5]) if whenever  $A, B \subset L$ ,  $|A|, |B| < \alpha$ ,  $B \geq A$ , and  $0 = \bigwedge (B - A)$ , then there is a  $c \in L$  such that  $c = \bigvee A = \bigwedge B$ . (We write  $B \geq A$  to mean  $b \geq a$  for all  $a \in A$ ,  $b \in B$ , and  $B - A$  for  $\{b - a: b \in B, a \in A\}$ .)

Recall  $L$  is called *Dedekind complete* if every subset of  $L$  that is bounded above has a supremum in  $L$ .  $L$  is  $\infty$ -Dedekind complete (as defined above) if and only if  $L$  is Dedekind complete. Two primary results of this paper are:

**Theorem 2.16.** Let  $\gamma: L \hookrightarrow M$  be an  $\alpha$ -jam-dense (see Proposition 2.5) embedding. Suppose  $\varphi: L \rightarrow N$  is  $\alpha$ -complete and  $N$  is  $\alpha$ -Dedekind complete. Then there exists a unique  $\alpha$ -complete morphism  $\bar{\varphi}: M \rightarrow N$  such that  $\varphi = \bar{\varphi} \circ \gamma$ .

**Theorem 2.19.** In  $\mathcal{W}(\alpha)$ , the full subcategory of  $\alpha$ -Dedekind complete objects is *epireflective*.

Other significant results, as we explain below, are the “topological realizations” of the two results above.

**Definition 1.3.**  $\text{Comp}$  denotes the category of compact Hausdorff spaces and continuous functions. In [2, 3, 15, 16], the contravariant *Yosida functor*,  $Y: \mathcal{W} \rightarrow \text{Comp}$  is investigated. For each  $L \in |\mathcal{W}|$ , there is a  $Y(L) \in |\text{Comp}|$ , called the Yosida space of  $L$  [32]; and for each  $\mathcal{W}$  morphism  $\varphi: L \rightarrow M$  there is a continuous function  $Y(\varphi): Y(M) \rightarrow Y(L)$ .

**Definition 1.4.** Let  $X, Y \in |\text{Comp}|$ . A continuous function  $f: X \rightarrow Y$  is called an  $\alpha$ -SpFi *morphism* if  $f^{-1}(K)$  is dense in  $X$  whenever  $K$  is dense and  $\alpha$ -Lindelöf in  $Y$ .  $\alpha$ -SpFi [4, 5, 20] denotes the topological category of compact Hausdorff spaces with  $\alpha$ -SpFi morphisms.

Below is a result which is essentially [3, 4.2]. It shows that  $Y$  restricted to  $\mathcal{W}(\alpha)$  is a functor to  $\alpha$ -SpFi.

**Theorem 3.10.** *A  $\mathcal{W}$  morphism  $\varphi: L \rightarrow M$  is  $\alpha$ -complete if and only if  $Y(\varphi): Y(M) \rightarrow Y(L)$  is an  $\alpha$ -SpFi morphism.*

Moreover,  $Y$  takes  $\mathcal{W}(\alpha)$  epics to  $\alpha$ -SpFi monics:

**Theorem 3.11.** *An  $\alpha$ -complete  $\mathcal{W}$  morphism  $\varphi: L \rightarrow M$  is epic in  $\mathcal{W}(\alpha)$  if and only if  $Y(\varphi): Y(M) \rightarrow Y(L)$  is monic in  $\alpha$ -SpFi.*

**Definition 1.5.** A Tychonoff topological space  $X$  is called  $\alpha$ -quasi- $F$  if each dense  $\alpha$ -Lindelöf set in  $X$  is  $C^*$ -embedded ( $\omega_1$ -quasi- $F \equiv$  quasi- $F$  [9, 18]).  $X$  is  $\infty$ -quasi- $F$  if and only if  $X$  is extremally disconnected [11, 28].

**Theorem 3.13.** *Let  $L \in |\mathcal{W}|$ . If  $L$  is  $\alpha$ -Dedekind complete, then  $Y(L)$  is  $\alpha$ -quasi- $F$ .*

The converse of this statement is not true; however:

**Theorem 3.12** [5].  *$X$  is  $\alpha$ -quasi- $F$  if and only if  $C(X)$  is  $\alpha$ -Dedekind complete.*

Applying  $Y$  to Theorem 2.19, we obtain the topological result:

**Theorem 4.3.** *In  $\alpha$ -SpFi, the full subcategory of  $\alpha$ -quasi- $F$  spaces is monoreflective.*

**Definition 1.6.** A *cover* of a compact space  $X$  is a pair  $(Y, f)$ , where  $Y$  is a compact space, and  $f$  maps  $Y$  irreducibly onto  $X$ . A continuous surjection  $f$  is *irreducible* if  $f$  does not map any proper closed subset of the domain onto the codomain. For an abstract topological property  $P$ , we say  $(Y, f)$  is a  *$P$  cover* of  $X$  if  $(Y, f)$  is a cover of  $X$ , and  $Y$  has the property  $P$ .  $(Y, f)$  is called the *minimum  $P$  cover* of  $X$  if given any other  $P$  cover  $(Z, g)$  of  $X$ , there is a unique irreducible map  $h: Z \rightarrow Y$  such that  $g = f \circ h$ .

In this paper (Theorem 4.3), we identify the  $\alpha$ -quasi- $F$  cover of a compact space  $X$ , denoted  $(QF_\alpha X, q_\alpha)$ , as the Yosida space of the  $\alpha$ -Dedekind complete epireflection, in  $\mathcal{W}(\alpha)$ , of  $C(X)$ . Moreover, in  $\alpha$ -SpFi,  $(QF_\alpha X, q_\alpha)$  is the  $\alpha$ -quasi- $F$  monoreflection of  $X$ .

## 2. The $\alpha$ -Dedekind complete epireflection

In this section, Zorn's lemma is used to obtain the  $\alpha$ -Dedekind complete epireflection of  $L$ .  $L$ ,  $M$ , and  $N$  denote  $\mathcal{W}$  objects, and maps between them are considered to be  $\mathcal{W}$  morphisms unless otherwise stated. The following is straightforward.

**Proposition 2.1.** *Let  $\varphi : L \rightarrow M$ . The following are equivalent.*

- (a)  $\varphi$  is  $\alpha$ -complete.
- (b) *There is a  $c \in L$  such that whenever  $B \subset L$ ,  $|B| < \alpha$ , and  $c = \bigvee^L B$ , then  $\varphi(c) = \bigvee^M \varphi[B]$ .*
- (c) *For  $A \subset L$  with  $|A| < \alpha$ , we have that  $\varphi(\bigwedge^L A) = \bigwedge^M \varphi[A]$  whenever  $\bigwedge^L A$  exists in  $L$ .*

**Proposition 2.2.** *Let  $\varphi : L \rightarrow M$  and  $\gamma : M \hookrightarrow N$ . If  $\gamma \circ \varphi$  is  $\alpha$ -complete, then  $\varphi$  is  $\alpha$ -complete.*

**Proof.** Let  $c \in L$  and suppose  $c = \bigvee^L A$  where  $A \subset L$  and  $|A| < \alpha$ . We claim that  $\varphi(c) = \bigvee^M \varphi[A]$ . Suppose not. Then there is a  $b \in M$  such that  $\varphi(c) > b > \varphi(a)$  for all  $a \in A$ . Because  $\gamma$  is injective, we have  $\gamma \circ \varphi(c) > \gamma(b) > \gamma \circ \varphi(a)$  for all  $a \in A$ . But this contradicts the assumption that  $\gamma \circ \varphi$  is  $\alpha$ -complete. For then  $\gamma \circ \varphi(c) = \bigvee^N \gamma \circ \varphi[A]$ .  $\square$

Henceforth,  $L \subseteq M$  denotes that  $L$  is a  $\mathcal{W}$  subspace of  $M$  (i.e.,  $L$  is a vector lattice subspace of  $M$ , and  $L$  has the same distinguished weak unit as  $M$ ).  $L \subseteq^\alpha M$  denotes that  $L$  is a  $\mathcal{W}$  subspace of  $M$ , and the inclusion of  $L$  into  $M$  is an  $\alpha$ -complete morphism. As usual, we reserve  $\subset$  for ordinary set inclusion.

**Definition 2.3.** Let  $L \subseteq M$ .  $L$  is said to be  $\alpha$ -dense in  $M$  if for each  $b \in M^+$  there is an  $A \subset L$  with  $|A| < \alpha$  such that  $b = \bigvee^M A$ .  $L$  is said to be  $\alpha$ -jam-dense in  $M$  if for each  $c \in M^+$  there are  $A, B \subset L$  with  $|A|, |B| < \alpha$  such that  $c = \bigwedge^M B = \bigvee^M A$ . (Or,  $0 = \bigwedge^M (B - A)$  and  $B \geq c \geq A$ .)

An embedding  $\gamma : L \hookrightarrow M$  is said to be  $\alpha$ -dense ( $\alpha$ -jam-dense) if  $\gamma[L]$  is  $\alpha$ -dense ( $\alpha$ -jam-dense) in  $M$ . Later we will see (Proposition 2.8) that  $\alpha$ -dense embeddings are epic in  $\mathcal{W}(\alpha)$ . Clearly,  $\alpha$ -jam-density implies  $\alpha$ -density, but the converse is not true. For  $C(X)$  is  $\omega_1$ -dense, but not  $\omega_1$ -jam-dense, in  $D(X)$  whenever  $D(X)$  is indeed a  $\mathcal{W}$  object. (See [17] and Section 3 here.)

**Proposition 2.4.** *Suppose  $L$  and  $M$  are  $\mathcal{S}$  objects. Then  $L$  is  $\alpha$ -dense in  $M$  if and only if  $L$  is  $\alpha$ -jam-dense in  $M$ .*

**Proof.** The sufficiency is clear. On the other hand, let  $L$  be  $\alpha$ -dense in  $M$ . For each  $c \in M^+$  there is  $A \subset L$  with  $|A| < \alpha$  such that  $c = \bigvee^M A$ . If  $u$  is the strong unit in  $L$  and  $M$ , there is an  $n \in \mathbb{N}$  such that  $nu > c$ . So  $nu - c \in M^+$ , and there is a  $B \subset L$  with  $|B| < \alpha$  such that  $nu - c = \bigvee^M B$ . It follows that  $c = \bigwedge^M \{nu - b : b \in B\}$ . Therefore  $L$  is  $\alpha$ -jam-dense in  $M$ .  $\square$

The following propositions are well known [6, 32, 26].

**Proposition 2.5.**  *$L$  is  $\infty$ -dense in  $M$  if and only if for each  $b \in M^+$  there is an  $a \in L^+$  such that  $b > a > 0$ .*

**Proposition 2.6.** *If  $L$  is  $\infty$ -dense in  $M$ , then  $L$  is  $\alpha$ -completely embedded in  $M$  for all  $\alpha < \infty$  (i.e.,  $L \subseteq^\alpha M$ ). Therefore if  $L$  is  $\alpha$ -dense in  $M$ , then  $L \subseteq^\alpha M$ .*

**Corollary 2.7.** *If  $\varphi : L \hookrightarrow M$  is  $\alpha$ -dense, then  $\varphi$  is  $\infty$ -complete (and hence  $\alpha$ -complete for all  $\alpha < \infty$ ).*

**Proof.**  $L$  is isomorphic to  $\varphi[L]$ , and  $\varphi[L] \subseteq^\infty M$  (Proposition 2.6). Therefore  $\varphi$  is  $\infty$ -complete because it is the composition of two  $\infty$ -complete morphisms.  $\square$

We may assume that  $L \subseteq^\alpha M$  whenever there is an  $\infty$ -dense embedding of  $L$  into  $M$ .

**Proposition 2.8.** *If  $\varphi : L \hookrightarrow M$  is  $\alpha$ -dense, then  $\varphi$  is epic in  $\mathcal{W}(\alpha)$ .*

**Proof.** By Corollary 2.7,  $\varphi$  is  $\alpha$ -complete and therefore a morphism in  $\mathcal{W}(\alpha)$ . Let  $\gamma_i : M \hookrightarrow N$ , with  $i = 1, 2$ , be  $\alpha$ -complete and suppose  $\gamma_1 \circ \varphi = \gamma_2 \circ \varphi$ . Let  $b \in M$ . Without loss of generality we may assume that  $b \in M^+$ . Therefore  $b = \bigvee^M \varphi[A]$  for some  $A \subset L$  with  $|A| < \alpha$ . Hence  $\gamma_1(b) = \gamma_1(\bigvee^M \varphi[A]) = \bigvee^N \gamma_1 \circ \varphi[A] = \bigvee^N \gamma_2 \circ \varphi[A] = \gamma_2(\bigvee^M \varphi[A]) = \gamma_2(b)$  because  $\gamma_1 \circ \varphi(a) = \gamma_2 \circ \varphi(a)$  for all  $a \in A$ .  $\square$

Proposition 2.2 says that if the second factor of an  $\alpha$ -complete morphism is injective, then the first factor of that  $\alpha$ -complete morphism is also  $\alpha$ -complete. Unfortunately the second factor of an  $\alpha$ -complete morphism need not be  $\alpha$ -complete; however:

**Lemma 2.9** (see [26, 23.2]). *Let  $\gamma : L \hookrightarrow M$  be  $\alpha$ -dense and  $\varphi : M \rightarrow N$ . If  $\varphi \circ \gamma$  is  $\alpha$ -complete, then  $\varphi$  is  $\alpha$ -complete.*

**Proof.** We may assume that  $L$  is  $\alpha$ -dense in  $M$ . Suppose  $\varphi$  is not  $\alpha$ -complete. Then by Proposition 2.1(b), there is for each  $c \in M^+$  a  $B_c \subset M^+$  with  $|B_c| < \alpha$  and  $c = \bigvee^M B_c$ , but  $\varphi(c) \neq \bigvee^N \varphi[B_c]$ . Therefore there is a  $d_c \in N^+$  such that  $\varphi(c) > d_c > \varphi(b)$  for all  $b \in B_c$ . Pick  $a \in L^+$  (hence  $a \in M^+$ ) and consider  $B_a$  as described above. For each  $b_i \in B_a$  there is an  $A_i \subset L$  with  $|A_i| < \alpha$  and  $b_i = \bigvee^M A_i$  because  $L$  is  $\alpha$ -dense in  $M$ . Moreover,  $a = \bigvee^M \bigcup_i A_i = \bigvee^L \bigcup_i A_i$  because  $a \in L^+$ , and  $\bigcup_i A_i \subset L$ . We claim that  $\varphi|_L$  is not  $\alpha$ -complete, i.e., we show that  $\varphi(a) \neq \bigvee^N \varphi[\bigcup_i A_i]$  ( $|\bigcup_i A_i| < \alpha$ ). As discussed above, there is a  $d_a \in N^+$  such that  $\varphi(a) > d_a > \varphi(b_i)$  for all  $b_i \in B_a$ . From this we conclude that  $\varphi(a) > d_a > \varphi(a_i)$  for all  $a_i \in A_i$  and for all  $i$ . Therefore  $\varphi(a) \neq \bigvee^N \varphi[\bigcup_i A_i]$ .  $\square$

Let  $L \subseteq M$  and  $b \in M^+$ . The smallest  $\mathcal{W}$  subspace of  $M$  that contains  $L$  and  $b$  is the smallest subset of  $M$  that contains  $L$  and  $b$  and is closed under the vector lattice operations. Using distributive laws in [6, 7, 10] we have:

**Proposition 2.10.** *Let  $L \subseteq M$  and  $b \in M$ . Then  $\langle L, b \rangle^M$ , the smallest  $\mathcal{W}$  subspace of  $M$  that contains  $L$  and  $b$ , is  $\{\bigwedge_{i=1}^n \bigvee_{j=1}^k a_{ij} + r_{ij}b : a_{ij} \in L, r_{ij} \in \mathbb{R}, n, k \in \mathbb{N}\}$ .*

Note, the superscript  $M$  in  $\langle L, b \rangle^M$  is omitted when the context is clear.

**Lemma 2.11.** *Let  $L \subseteq M$ ,  $A, B \subset L$ ,  $|A|, |B| < \alpha$ ,  $b \in M^+$ , and  $b = \bigvee^M A = \bigwedge^M B$ . Then  $L$  is  $\alpha$ -jam-dense in  $\langle L, b \rangle$ .*

**Proof.** Let  $c \in \langle L, b \rangle^+$ . Then  $c = \bigwedge_{i=1}^n \bigvee_{j=1}^k (a_{ij} + r_{ij}b)$  for some  $a_{ij} \in L$ ,  $r_{ij} \in \mathbb{R}$ , and  $n, k \in \mathbb{N}$ . For each  $(i, j)$  let  $d_{ij} = a_{ij} + r_{ij}b = a_{ij} + r_{ij} \bigvee^M A = a_{ij} + r_{ij} \bigwedge^M B$ . We have

$$c = \bigwedge_{i=1}^n \bigvee_{j=1}^k d_{ij}.$$

Now for  $r_{ij} > 0$ ,

$$d_{ij} = \bigvee^M \{a_{ij} + r_{ij}a : a \in A\} = \bigwedge^M \{a_{ij} + r_{ij}p : p \in B\},$$

and for  $r_{ij} \leq 0$ ,

$$d_{ij} = \bigwedge^M \{a_{ij} + r_{ij}a : a \in A\} = \bigvee^M \{a_{ij} + r_{ij}p : p \in B\}.$$

For each  $(i, j)$  define the sets

$$C_{ij} = \begin{cases} \{a_{ij} + r_{ij}a : a \in A\}, & \text{if } r_{ij} > 0, \\ \{a_{ij} + r_{ij}p : p \in B\}, & \text{if } r_{ij} \leq 0, \end{cases}$$

$$D_{ij} = \begin{cases} \{a_{ij} + r_{ij}p : p \in B\}, & \text{if } r_{ij} > 0, \\ \{a_{ij} + r_{ij}a : a \in A\}, & \text{if } r_{ij} \leq 0. \end{cases}$$

Then  $d_{ij} = \bigvee^M C_{ij} = \bigwedge^M D_{ij}$ .

It follows that  $c = \bigwedge_{i=1}^n \bigvee_{j=1}^k (\bigvee^M C_{ij}) = \bigwedge_{i=1}^n \bigvee_{j=1}^k (\bigwedge^M D_{ij})$ , and  $c = \bigvee^M C' = \bigwedge^M D'$  with  $C', D' \subset L$  and  $|C'|, |D'| < \alpha$ . Since  $c \in \langle L, b \rangle$ ,  $c = \bigvee^{\langle L, b \rangle} C' = \bigwedge^{\langle L, b \rangle} D'$ . Hence  $L$  is  $\alpha$ -jam-dense in  $\langle L, b \rangle$ .  $\square$

In light of Lemma 2.11 one may ask: if  $L \subseteq M$ ,  $b \in M^+$ ,  $A \subset L$ ,  $|A| < \alpha$ , and  $b = \bigvee^M A$ , then is  $L$   $\alpha$ -dense in  $\langle L, b \rangle$ ? The answer is “no” (Example 2.12) even if both  $L$  and  $M$  are  $\mathcal{S}$  objects.

**Example 2.12.** Let  $X \in |\text{Comp}|$  be extremally disconnected [11]. Suppose  $U$  is open, but not clopen in  $X$ . Let  $L(X)$  be the  $\mathcal{W}$  object of locally constant real-valued functions on  $X$  [10, 32]. Consider  $L(X) \subseteq \mathbb{R}^X$  ( $\mathbf{1}$  is the strong unit). The characteristics function of  $U$ ,  $\chi_U$ , is not an element of  $L(X)$ , and clearly  $\chi_U = \bigvee^{\mathbb{R}^X} \{b \in L(X) : b < \chi_U\}$ . (Actually,  $\chi_U$  is the sup of a subset of  $L(X)$  whose cardinality is equal to the weight of  $X$ .) We claim that  $L(X)$  is not  $\infty$ -dense in  $\langle L(X), \chi_U \rangle$ .

Let  $K = \bar{U} - U$ . Since  $1 - \chi_U = \chi_{X \setminus U} \in \langle L(X), \chi_U \rangle^+$ ,  $\chi_K = \chi_{X \setminus U} \wedge \chi_{\bar{U}} \in \langle L(X), \chi_U \rangle^+$ . However, since  $K$  is nowhere dense, there is no  $b \in L(X)^+$  such that  $\chi_K > b > 0$ . Therefore, by Proposition 2.5,  $L(X)$  is not  $\infty$ -dense in  $\langle L(X), \chi_U \rangle$ .

If  $L \subseteq N \subseteq M$  and  $L$  is  $\alpha$ -dense ( $\alpha$ -jam-dense) in  $N$ ,  $N$  is an  $\alpha$ -dense ( $\alpha$ -jam-dense) extension of  $L$  in  $M$ . Below we show there are maximal, sometimes maximum, such extensions.

**Proposition 2.13.** *Let  $L \subseteq M$  and  $P$  be a family of subspaces of  $M$ . View  $P$  as a poset ordered by set inclusion and suppose  $C$  is a chain in  $P$ .*

- (a) *If  $P = \{L_i \subseteq M : L \subseteq^\alpha L_i\}$ , then  $\bigcup C \in P$ , i.e.,  $L \subseteq^\alpha \bigcup C$ .*
- (b) *If  $P = \{L_i \subseteq M : L \text{ is } \alpha\text{-dense in } L_i\}$ , then  $\bigcup C \in P$ , i.e.,  $L$  is  $\alpha$ -dense in  $\bigcup C$ .*
- (c) *If  $P = \{L_i \subseteq M : L \text{ is } \alpha\text{-jam-dense in } L_i\}$ , then  $\bigcup C \in P$ , i.e.,  $L$  is  $\alpha$ -jam-dense in  $\bigcup C$ .*

**Proof.** Obviously an increasing union of  $\mathcal{W}$  subobjects in  $M$  is a  $\mathcal{W}$  subobject of  $M$ .

(a) Suppose  $L$  is not  $\alpha$ -completely embedded in  $\bigcup C$ . Then there is a  $c \in L_i^+ \in C$  and an  $A \subset L$  with  $|A| < \alpha$  and  $c = \bigvee^{L_i} A$  such that  $c \neq \bigvee^{\bigcup C} A$ . So there is a  $b \in \bigcup C$  such that  $c > b > a$  for all  $a \in A$ . Since  $b \in L_i$  for some  $L_i \in C$ ,  $c \neq \bigvee^{L_i} A$ . But this is a contradiction since  $L \subseteq^\alpha L_i$ .

(b) Since  $L \subseteq^\alpha L_i$  (Proposition 2.6) we have by (a) here that  $L \subseteq^\alpha \bigcup C$ . Let  $b \in \bigcup C^+$ . Then  $b \in L_i^+ \in C$  for some  $i$ , and there is an  $A \subset L$  with  $|A| < \alpha$  and  $b = \bigvee^{L_i} A$ . By Lemma 2.9,  $L_i \subseteq^\alpha \bigcup C$ , hence  $b = \bigvee^{\bigcup C} A$ . Therefore  $L$  is  $\alpha$ -dense in  $\bigcup C$ .

(c) As in (b),  $L \subseteq^\alpha \bigcup C$ . Let  $c \in \bigcup C^+$ . Then  $c \in L_i^+ \in C$  for some  $i$ , and there are  $A, B \subset L$  with  $|A|, |B| < \alpha$  such that  $c = \bigwedge^{L_i} B = \bigvee^{L_i} A$ . By Lemma 2.9,  $L_i \subseteq^\alpha \bigcup C$ , and it follows from Proposition 2.1 that  $c = \bigvee^{\bigcup C} A = \bigwedge^{\bigcup C} B$  and  $L$  is  $\alpha$ -jam-dense in  $\bigcup C$ .  $\square$

**Lemma 2.14.** *Let  $L \subseteq M$ . There are  $L_1, L_2$ , and  $L_3$  with  $L \subseteq^\alpha L_i \subseteq M$  such that:*

- (a)  *$L_1$  is maximal for the property of containing  $L$  as an  $\alpha$ -completely embedded subspace.*
- (b)  *$L_2$  is maximal for the property of containing  $L$  as an  $\alpha$ -dense subspace.*
- (c)  *$L_3$  is maximal for the property of containing  $L$  as an  $\alpha$ -jam-dense subspace.*

**Proof.** Proposition 2.13 allows us to invoke Zorn's lemma in each of the above cases.  $\square$

If in Lemma 2.14,  $L \subseteq^\alpha M$ , we have:

**Theorem 2.15.** *Let  $L \subseteq^\alpha M$ . There is a subspace of  $M$ , denoted  $[L]_\alpha^M$ , such that  $[L]_\alpha^M$  is maximum (among the subspaces of  $M$ ) for the property of containing  $L$  as an  $\alpha$ -jam-dense subspace. If  $L$  and  $M$  are  $\mathcal{S}$  objects, then  $[L]_\alpha^M$  is maximum for the property of containing  $L$  as an  $\alpha$ -dense subspace (Proposition 2.4). Moreover, if  $M$  is  $\alpha$ -Dedekind complete, then  $[L]_\alpha^M$  is also  $\alpha$ -Dedekind complete.*

**Proof.** Suppose  $L_1, L_2, L_1 \neq L_2$ , are maximal for the property of containing  $L$  as an  $\alpha$ -jam-subspace. Then there is a  $b \in L_1^+ \setminus L_2^+$ , and there are  $A, B \subset L_1$  with  $|A|, |B| < \alpha$  such that  $b = \bigvee^{L_1} A = \bigwedge^{L_1} B$ . Since  $L_1 \subseteq^\alpha M$  (Proposition 2.6),  $b = \bigvee^M A = \bigwedge^M B$ . Then by Example 2.12,  $L_2$  is  $\alpha$ -jam-dense in  $\langle L_2, b \rangle$ . But this contradicts the maximality of  $L_2$ .

Now suppose that  $M$  is  $\alpha$ -Dedekind complete, but  $[L]_\alpha^M$  is not. Then there are  $A, B \subset [L]_\alpha^{M+}$  with  $|A| < \alpha, |B| < \alpha, B > A$ , and  $0 = \bigwedge^{[L]_\alpha^M} (B - A)$  for which there does not exist a  $c \in [L]_\alpha^M$  satisfying  $B > c > A$ .  $[L]_\alpha^M \subseteq^\alpha M$ , so  $0 = \bigwedge^M (B - A)$  (Proposition 2.1), and since  $M$  is  $\alpha$ -Dedekind complete, there is a  $c \in M$  such that  $c = \bigvee^M A = \bigwedge^M B$ . Hence by Lemma 2.11,  $[L]_\alpha^M$  is  $\alpha$ -jam-dense in  $\langle [L]_\alpha^M, c \rangle$ , but this contradicts the maximality of  $[L]_\alpha^M$ .  $\square$

We now give the main result of this section.

**Theorem 2.16.** *Let  $\gamma: L \hookrightarrow M$  be an  $\alpha$ -jam-dense embedding and suppose  $\varphi: L \rightarrow N$  is  $\alpha$ -complete with  $N$   $\alpha$ -Dedekind complete. Then there is a unique  $\bar{\varphi}: M \rightarrow N$  such that  $\bar{\varphi} \circ \gamma = \varphi$ .  $\bar{\varphi}$  is necessarily  $\alpha$ -complete. Moreover, if  $L$  and  $M$  are  $\mathcal{S}$  objects, then the same lifting result holds when  $\gamma$  is an  $\alpha$ -dense embedding (Proposition 2.4).*

**Proof.** We may assume that  $L$  is an  $\alpha$ -jam-dense subspace of  $M$ . Also, if  $\bar{\varphi}$  exists, it is  $\alpha$ -complete (Lemma 2.9), and hence unique (Proposition 2.8).

We define  $\bar{\varphi}$ . Let  $b \in M$ . Without loss of generality we can assume that  $b \in M^+$ . There are  $A, B \subset L$  with  $|A|, |B| < \alpha$ , with  $b = \bigvee^M A = \bigwedge^M B$ . So  $0 = \bigwedge^M (B - A)$ . Therefore by Proposition 2.1,  $0 = \varphi(0) = \bigwedge^N (\varphi[B] - \varphi[A])$ . From this it follows, since  $N$  is  $\alpha$ -Dedekind complete, that there is a  $c \in N$  such that  $c = \bigvee^N \varphi[A] = \bigwedge^N \varphi[B]$ . Define  $\bar{\varphi}(b) = c$ . Clearly  $\bar{\varphi}$  is an assignment that extends  $\varphi$ , and it is not hard to see that  $\bar{\varphi}$  is a  $\mathcal{W}$  morphism if, as is shown below, it is well defined.

Let  $d \in M$  and suppose  $d = \bigvee^M A_1 = \bigwedge^M B_1 = \bigvee^M A_2 = \bigwedge^M B_2$  for  $A_i, B_i$  as above. It follows that there are  $c_1, c_2 \in N$  such that

$$c_1 = \bigvee^N \varphi[A_1] = \bigwedge^N \varphi[B_1], \quad \text{and} \quad c_2 = \bigvee^N \varphi[A_2] = \bigwedge^N \varphi[B_2].$$

To see that  $\bar{\varphi}$  is well defined it suffices to see that  $c_1 = c_2$ .

To this end, we claim that  $c_1 \geq \varphi(a')$  for all  $a' \in A_2$ , and  $c_1 \leq \varphi(b')$  for all  $b' \in B_2$ . From this it follows that  $c_1 = c_2 = \bigvee^N \varphi[A_2] = \bigwedge^N \varphi[B_2]$ .

For each  $a' \in A_2 \subset L$  we have

$$\bigvee^M \{a' \wedge a: a \in A_1\} = a' \wedge \bigvee^M A_1 = a' \wedge d = a'.$$

Therefore  $\bigvee^L \{a' \wedge a: a \in A_1\} = a'$ . Since  $\varphi$  is  $\alpha$ -complete, we have that

$$\begin{aligned} \varphi(a') &= \bigvee^N \{\varphi(a') \wedge \varphi(a): a \in A_1\} \\ &= \varphi(a') \wedge \bigvee^N \varphi[A_1] = \varphi(a') \wedge c_1. \end{aligned}$$



Therefore  $c_1 \geq \varphi(a')$  for all  $a' \in A_2$ . Similarly for each  $b' \in B_2 \subset L$  we have,

$$\bigwedge^M \{b' \vee b : b \in B_1\} = b' \vee \bigwedge^M B_1 = b' \vee d = b'.$$

Hence  $\bigwedge^L \{b' \vee b : b \in B_1\} = b'$ . It follows that

$$\begin{aligned} \varphi(b') &= \bigwedge^N \{\varphi(b') \vee \varphi(b) : b \in B_1\} \\ &= \varphi(b') \vee \bigwedge^N \varphi[B_1] = \varphi(b') \vee c_1. \end{aligned}$$

Therefore  $\varphi(b') \geq c_1$  for all  $b' \in B_2$ .  $\square$

The next theorem is well known and it tells us that each  $L$  has an essentially unique Dedekind ( $\infty$ -Dedekind complete) completion [32].

**Theorem 2.17.** *Each  $L$  has an essentially unique pair  $(\sigma, \bar{L})$  where  $\bar{L}$  is Dedekind complete and  $\sigma$  is an  $\infty$ -jam-dense embedding. That is, for each  $b \in \bar{L}^+$ ,*

$$b = \bigvee^{\bar{L}} \{\sigma(a) : \sigma(a) \leq b\} = \bigwedge^{\bar{L}} \{\sigma(a) : \sigma(a) \geq b\}.$$

$(\sigma, \bar{L})$  is called the Dedekind completion of  $L$ .

When we say  $(\sigma, \bar{L})$  is *essentially unique*, we mean that if given another pair  $(\sigma', \bar{L}')$  with the same properties (as in Theorem 2.17), there is an isomorphism,  $\tau: \bar{L} \rightarrow \bar{L}'$ , such that  $\sigma' = \tau \circ \sigma$ . Because of this, we will suppress reference to the embedding  $\sigma$ . That is, we can consider  $L$  to be an  $\infty$ -jam-dense subspace of a unique Dedekind complete  $\mathcal{W}$  object  $\bar{L}$ ; and we say that  $\bar{L}$  “is” the Dedekind completion of  $L$ . We begin to generalize Theorem 2.17 to arbitrary  $\alpha$ . For an  $L$ , we say that  $(\varphi, M)$  is an  $\alpha$ -jam-dense ( $\alpha$ -dense) extension of  $L$  if  $\varphi: L \hookrightarrow M$  is  $\alpha$ -jam-dense ( $\alpha$ -dense).

**Proposition 2.18.** *An  $\alpha$ -jam-dense extension,  $(\varphi, M)$ , of  $L$ , with  $M$   $\alpha$ -Dedekind complete is essentially unique.*

**Proof.** Apply Theorem 2.16.  $\square$

Recall in an abstract category  $\mathcal{B}$ , a full subcategory  $\mathcal{A}$  is called *epireflective* if for each  $B \in |\mathcal{B}|$  there is an  $A_B \in |\mathcal{A}|$  and an epimorphism  $e: B \rightarrow A_B$  such that for any morphism  $f: B \rightarrow A$  with  $A \in |\mathcal{A}|$  there exists an  $\bar{f}: A_B \rightarrow A$  such that  $f = \bar{f} \circ e$ .  $(e, A_B)$  is called the  $\mathcal{A}$  *epireflection* of  $B$ .

We claim that, in  $\mathcal{W}(\alpha)$ , the full subcategory of  $\alpha$ -Dedekind complete objects is epireflective. Because of the lifting theorem (Theorem 2.16), we need only produce, for each  $L$ , an  $\alpha$ -jam-dense extension, call it  $(i, \hat{L}_\alpha)$ , with  $\hat{L}_\alpha$   $\alpha$ -Dedekind complete. This is easily done. Recall that  $L$  is  $\infty$ -jam-dense in  $\bar{L}$ . Define  $\hat{L}_\alpha = [L]_\alpha^{\bar{L}}$  to be the maximum  $\alpha$ -jam-dense extension of  $L$  in  $\bar{L}$  (Theorem 2.15).  $\hat{L}_\alpha$  is  $\alpha$ -Dedekind complete because  $\bar{L}$  is  $\alpha$ -Dedekind complete for all  $\alpha < \infty$ . Hence  $(i, \hat{L}_\alpha)$  is the  $\alpha$ -Dedekind complete epireflection of  $L$  where  $i$  is the inclusion of  $L$  into  $\hat{L}_\alpha$ . ( $i$  is epic by Proposition 2.8.) Note,  $\hat{L}_\infty = \bar{L}$ . We summarize below.

**Theorem 2.19.** *In  $\mathcal{W}(\alpha)$ , the full subcategory of  $\alpha$ -Dedekind complete objects is epireflective, and for each  $L$ ,  $(i, \hat{L}_\alpha)$  is the  $\alpha$ -Dedekind complete epireflection of  $L$ .*

### 3. The Yosida functor

In this section,  $X, Y, Z \in |\mathbf{Comp}|$ ;  $f, g$ , and  $h$  denote continuous functions.

The functor  $Y$  works very much like the Stone functor from Boolean algebras to Boolean spaces.  $Y(L)$ , like the Stone space of a Boolean algebra, is a maximal ideal space. The elements of  $Y(L)$  are ideals of  $L$  that are maximal for the property of not containing the weak unit. If the weak unit is a strong unit, then these ideals are the actual maximal ideals of  $L$ . The topology on this space is the hull-kernel topology. In fact, if we view a Boolean algebra,  $B$ , as an  $\mathcal{S}$  object (i.e., consider  $L(S(B))$ ), the locally constant real-valued functions on the Stone space,  $S(B)$ , of  $B$ , the Yosida functor can be thought of as an extension of the Stone functor.

The following discussion comes from [2]. Let  $C(X)$  be the  $\mathcal{S}$  object of real-valued continuous functions on  $X$ . The strong unit of  $C(X)$  will always be taken to be the constant function  $\mathbf{1}$ . Let  $D(X)$  be the set of extended real-valued continuous functions,  $f: X \rightarrow [-\infty, +\infty]$ , for which  $f^{-1}(\mathbb{R})$  is dense in  $X$ . In the pointwise order,  $D(X)$  is a lattice, but usually fails to be a vector space. For  $f, g, h \in D(X)$ , we say “ $f + g = h$  in  $D(X)$ ” if  $f(x) + g(x) = h(x)$  when  $x \in f^{-1}(\mathbb{R}) \cap g^{-1}(\mathbb{R}) \cap h^{-1}(\mathbb{R})$  (which is a dense set in  $X$ ). It may well happen that, for particular  $f, g \in D(X)$ , there is no  $h \in D(X)$  with  $f + g = h$  in  $D(X)$  (e.g., take  $X = [-\infty, +\infty]$ ,  $f$  the obvious extension of  $x + \sin x$ , and  $g$  the extension of  $-x$ ). However, it may well happen that a subset  $L \subset D(X)$  has the property that for all  $f, g \in L$  there is an  $h \in D(X)$  with  $f + g = h$  in  $D(X)$ ; if  $L$  is also a vector lattice under the pointwise operation in  $D(X)$  and the constant function,  $\mathbf{1}$ , is in  $L$ , then we say “ $(L, \mathbf{1})$  (or just  $L$ ) is a  $\mathcal{W}$  object in  $D(X)$ ” (e.g.,  $C(X)$  is a vector lattice, indeed, a  $\mathcal{W}$  object, in  $D(X)$ ). If  $X$  has the property that each dense cozero set is  $C^*$ -embedded [11], then  $X$  is called  $\omega_1$ -quasi- $F$  (or just quasi- $F$ ). See Definition 1.5 here and [5, 9, 18]. If  $X$  is  $\omega_1$ -quasi- $F$ , then  $(D(X), \mathbf{1}) \in |\mathcal{W}|$ . See [17].

**Theorem 3.1** [15]. (a) *There is a  $\mathcal{W}$  isomorphism,  $\hat{\cdot}: L \rightarrow \hat{L} \subset D(Y(L))$ , onto a  $\mathcal{W}$  object,  $\hat{L}$  in  $D(Y(L))$ , with weak unit  $\hat{w}_L = \mathbf{1}$ , and  $\hat{L}$  separates the points of  $Y(L)$ .*

(b) *If  $L'$  is a  $\mathcal{W}$  object in  $D(X)$  which separates the points of  $X$ , and for  $a \in L$ , if  $a \mapsto a'$  is a  $\mathcal{W}$  isomorphism from  $L$  to  $L'$ , then there is a homeomorphism  $f: X \rightarrow Y(L)$  such that  $a' = \hat{a} \circ f$  for all  $a \in L$ .*

Theorem 3.1(b) is used to recognize Yosida representations.

Let  $L^* = (w_L)$  be the principal ideal generated by  $w_L$  in  $L$ .  $\hat{L}^*$  is a  $\mathcal{W}$  object (indeed, an  $\mathcal{S}$  object) in  $D(Y(L))$  and consists of all  $\hat{a} \in \hat{L}$  which are bounded. Note that if  $L$  is an  $\mathcal{S}$  object, i.e., if  $w_L$  is a strong unit, then  $L = L^*$ .

- Corollary 3.2.** (a)  $Y(L^*) = Y(L)$ .  
 (b)  $Y(C(X)) = X$ .  
 (c)  $\hat{L}^*$  is an  $\mathcal{S}$  subspace of  $C(Y(L))$ .  
 (d) If  $L$  is an  $\mathcal{S}$  object, then  $\hat{L} \subseteq C(Y(L))$ .

- Proof.** (a) In  $D(Y(L))$ ,  $\hat{L}^*$  satisfies Theorem 3.1(b).  
 (b) In  $D(X)$ ,  $C(X)$  satisfies Theorem 3.1(b).  
 (c)  $\hat{a} \in \hat{L}^*$  implies  $\hat{a}^{-1}(\{\pm\infty\}) = \emptyset$ , hence  $\hat{a} \in C(Y(L))$ .  
 (d) If  $L \in |\mathcal{S}|$ , then  $L = L^*$ .  $\square$

Let us look at a couple of simple examples. Since  $\mathbb{R}^X = C(D_X)$ , where  $D_X$  is the set  $X$  with the discrete topology, we can see from Corollary 3.2(a), (b) that  $Y(\mathbb{R}^X) = \beta D_X$  ( $C^*(D_X) = C(\beta D_X)$ ).

Let  $X$  be zero dimensional. Since  $L(X) \subseteq C(X)$  and  $L(X)$  (Example 2.12) separates the points of  $X$ , we can see from Theorem 3.1(b) and Corollary 3.2(b) that  $Y(L(X)) = X$ .

**Theorem 3.3** [15]. Let  $\varphi_i: L \rightarrow M$  for  $i = 1, 2$ .

- (a) There is a unique continuous function,  $Y(\varphi_1): Y(M) \rightarrow Y(L)$ , such that  $\varphi(a)^\wedge = \hat{a} \circ Y(\varphi_1)$  for all  $a \in L$ .  
 (b)  $Y$  is a faithful functor, i.e., if  $\varphi_1 \neq \varphi_2$ , then  $Y(\varphi_1) \neq Y(\varphi_2)$ .  
 (c)  $\varphi_1$  is one-to-one if and only if  $Y(\varphi_1)$  is onto, and if  $\varphi_1$  is onto, then  $Y(\varphi_1)$  is one-to-one.

Henceforth,  $L$  and  $\hat{L}$  are identified.

We will consider  $a \in L$  as an extended real-valued function on  $Y(L)$ , and if  $L$  is an  $\mathcal{S}$  object, we will consider  $a \in L$  as a real-valued function on  $Y(L)$ . See Corollary 3.2. In general,  $Y(\varphi)$  being one-to-one does not imply that  $\varphi$  is onto: if  $\varphi$  is the inclusion of  $L(X)$  (Example 2.12) into  $C(X)$ , then  $Y(\varphi): X = Y(C(X)) \rightarrow Y(L(X)) = X$  is  $\text{id}_X$ . However:

**Proposition 3.4.** Let  $\varphi: C(X) \rightarrow M$  and  $M \in |\mathcal{S}|$ . Then  $Y(\varphi): Y(M) \rightarrow X$  is one-to-one if and only if  $\varphi$  is onto.

**Proof.** For the sufficiency apply Theorem 3.3(c).

On the other hand, if  $Y(\varphi)$  is one-to-one, then  $Y(M)$  is homeomorphic (via  $Y(\varphi)$ ) to a closed subspace  $K$  of  $X$ . Hence there is a homeomorphism,  $Y(\varphi)^{-1}: K \rightarrow Y(M)$ . Let  $b \in M$ . Since  $M \subseteq C(Y(M))$ , we have that  $b \circ Y(\varphi)^{-1} \in C(K)$ . Therefore, there is a continuous extension,  $\bar{b} \in C(X)$ , of  $b \circ Y(\varphi)^{-1}$  to all of  $X$ . Then, by Theorem 3.3(a),  $\varphi(\bar{b}) = \bar{b} \circ Y(\varphi) = b$ .  $\square$

The next proposition is straightforward.

**Proposition 3.5.** *Let  $f: X \rightarrow Y$ . Define  $f': C(Y) \rightarrow C(X)$  by  $f'(g) = g \circ f$  for  $g \in C(Y)$  [11, Ch. 10]. Then  $f'$  is a  $\mathcal{W}$  morphism and  $Y(f') = f$ .*

**Theorem 3.6** (Banach-Stone). *A function  $f: X \rightarrow Y$  is a homeomorphism if and only if  $f': C(Y) \rightarrow C(X)$  is an isomorphism in  $\mathcal{W}$ .*

**Proof.**  $Y(f') = f$ . Apply Theorem 3.3(c) and Proposition 3.4.  $\square$

We begin a discussion of the topological category  $\alpha\text{-SpFi}$ . See Definition 1.4 here, and [4, 5, 20]. A topological space,  $K$ , is called  $\alpha$ -Lindelöf if any open cover of  $K$  has a subcover of cardinality less than  $\alpha$ . A subset of  $X$  is called an  $F_\alpha$ -set in  $X$  if it is the union of fewer than  $\alpha$  many closed subsets of  $X$ . Clearly every  $F_\alpha$ -set in  $X$  is  $\alpha$ -Lindelöf.

**Definition 3.7.** Let  $\text{Coz}(X) = \{f^{-1}(\mathbb{R} \setminus \{0\}) : f \in C(X)\}$ . A subset  $V \subset X$  is said to be an  $\alpha$ -cozero set if  $V = \bigcup \{U_i : i \in I, |I| < \alpha, U_i \in \text{Coz}(X)\}$ . Note that an  $\omega_1$ -cozero set is a cozero set. Recall that by “ $|I| < \infty$ ” we mean that “ $|I|$  is unrestricted”, so that every open set is an  $\infty$ -cozero set. We denote the collection of  $\alpha$ -cozero sets of  $X$  by  $\text{Coz}_\alpha(X)$ . Let  $\mathcal{G}_\alpha(X)$  be the filter generated by the dense members of  $\text{Coz}_\alpha(X)$ .  $\mathcal{G}_\infty(X)$  will denote the filter generated by the dense open sets. Finally let  $\mathcal{G}_\alpha(X)_\delta$  be the filter generated by

$$\{\bigcap G_n : n \in \mathbb{N}, G_n \in \mathcal{G}_\alpha(X)\}.$$

Note the Baire category theorem implies that  $\mathcal{G}_\alpha(X)_\delta$  is a filter of dense sets. Moreover, when  $\omega_0$  is replaced by  $\alpha$  [8, 9.7, 9.8, and 9.10(a)] show that the members of  $\mathcal{G}_\alpha(X)_\delta$  are  $\alpha$ -Lindelöf.

$\alpha\text{-SpFi}$  is actually a full subcategory of the more general category of *spaces with filters*, denoted  $\text{SpFi}$ , which has for objects pairs of the form  $(X, \mathcal{F})$  where  $\mathcal{F}$  is a filter base of dense subsets of  $X$ . A  $\text{SpFi}$  morphism  $f: (X, \mathcal{F}) \rightarrow (Y, \mathcal{H})$  is a continuous function from  $X$  to  $Y$  that inversely preserves the elements of the filter bases, i.e.,  $f^{-1}(H) \in \mathcal{F}$  for all  $H \in \mathcal{H}$ .

**Proposition 3.8.** *Let  $f: X \rightarrow Y$ . The following are equivalent.*

- (a)  *$f$  is an  $\alpha\text{-SpFi}$  morphism.*
- (b)  *$f^{-1}(G)$  is dense in  $X$  for each  $G \in \mathcal{G}_\alpha(Y)$ , i.e.,  $f: (X, \mathcal{G}_\alpha(X)) \rightarrow (Y, \mathcal{G}_\alpha(Y))$  is a  $\text{SpFi}$  morphism.*
- (c)  *$f^{-1}(H)$  is dense in  $X$  for each  $H \in \mathcal{G}_\alpha(Y)_\delta$ , i.e.,  $f: (X, \mathcal{G}_\alpha(X)_\delta) \rightarrow (Y, \mathcal{G}_\alpha(Y)_\delta)$  is a  $\text{SpFi}$  morphism.*

**Proof.** The equivalence of (b) and (c) is clear. Also (a) $\Rightarrow$ (b) is trivial because each  $G \in \mathcal{G}_\alpha(Y)$  is an  $F_\alpha$ -set in  $Y$  and hence  $\alpha$ -Lindelöf. For (b) $\Rightarrow$ (a) we use [4, 1.6]: if  $f$  is as in (b),  $U$  is regular closed in  $X$ , and  $G \in \mathcal{G}_\alpha(Y)$ , then  $f[U] \cap G$  is dense in  $f[U]$ .

So now let  $K$  be dense and  $\alpha$ -Lindelöf in  $Y$  and suppose  $f^{-1}(K)$  is not dense in  $X$ . Then there is a nonempty regular closed set  $U$  in  $X$  such that  $U \cap f^{-1}(K) = \emptyset$ . Therefore  $f[U] \cap K = \emptyset$ . It follows that there exists an  $\alpha$ -cozero set  $V$  in  $Y$  such that  $K \subset V$  and  $V \cap f[U] = \emptyset$ . But this contradicts [4, 1.6].  $\square$

The next three lemmas (Lemmas 3.9–3.11) provide a fundamental link between  $\mathcal{W}(\alpha)$  and  $\alpha$ -SpFi. (In regard to Lemmas 3.9 and 3.10, the essential ideas and content come from [3, 4.1 and 4.2].)

**Lemma 3.9.** Let  $b \in L^+$ ,  $A \subset L$ ,  $|A| < \alpha$ ,  $r \in \mathbb{R}$  and let  $P_b = \{x \in Y(L) : b(x) = \bigvee^{\mathbb{R}} \{a(x) : a \in A\}\}$ .

- (a)  $b = \bigvee^L A$  if and only if  $P_b \in \mathcal{G}_\alpha(Y(L))_\delta$ .
- (b) If  $r = \bigvee^L A$ , then  $\bigcup \{\text{coz}(a) : a \in A\} \in \mathcal{G}_\alpha(Y(L))$ .
- (c) If  $G \in \mathcal{G}_\alpha(Y(L))$ , then for each  $r \in \mathbb{R}$  there is a family  $A_r \subset L^+$  with  $|A_r| < \alpha$  such that  $G = \bigcup \{\text{coz}(a) : a \in A_r\}$  and  $r = \bigvee^L A_r$ .

**Proof.** (a) If  $P_b \in \mathcal{G}_\alpha(Y(L))_\delta$ , then  $P_b$  is dense in  $Y(L)$ , hence  $b = \bigvee^L A$ . On the other hand, suppose  $b = \bigvee^L A$ . For each  $n \in \mathbb{N}$  let  $U_n = \{x \in Y(L) : b(x) - a(x) < n^{-1} \text{ for some } a \in A\}$ . It is not hard to see that each  $U_n \in \text{Coz}_\alpha(Y(L))$  and  $P_b = \bigcap_n U_n$ . If each  $U_n$  is dense in  $Y(L)$ , we are finished. Suppose not, then there is an open set  $V \subset Y(L)$  for which  $b(x) - a(x) \geq n^{-1}$  for all  $x \in V$  and for all  $a \in A$ . Take another open set,  $V' \subset V$ , with  $\bar{V}' \subset V$ . Since  $L$  separates the points of  $Y(L)$  (Theorem 3.1), there is a  $c \in L^+$  such that  $c[Y(L) \setminus V] = 0$ ,  $c[\bar{V}'] = n^{-1}$ , and  $0 \leq c \leq n^{-1}$ . It follows that  $b - c > a$  for all  $a \in A$ , but this contradicts  $b = \bigvee^L A$ .

(b) If  $r = \bigvee^L A$ , then by (a),  $P_r$  is dense in  $Y(L)$ . Since  $P_r \subset \bigcup \{\text{coz}(a) : a \in A\}$ , it follows that  $\bigcup \{\text{coz}(a) : a \in A\} \in \mathcal{G}_\alpha(Y(L))$ .

(c) Let  $G \in \mathcal{G}_\alpha(Y(L))$  and  $x \in G$ . There is an  $a_x \in L^+$ , and a neighborhood  $V_x$  of  $x$  such that  $\bar{V}_x \subset G$ ,  $a_x[\bar{V}_x] = r$ ,  $a_x[X \setminus G] = 0$ , and  $0 < a_x \leq r$ . Since  $G$  is  $\alpha$ -Lindelöf, there is a subfamily  $\{V_{x'}\}$  of cardinality less than  $\alpha$  with  $G = \bigcup \{V_{x'}\}$ . Therefore since  $V_{x'} \subset \text{coz}(a_{x'}) \subset G$ , we have  $G = \bigcup \{\text{coz}(a_{x'})\}$ . Clearly,  $r = \bigvee^L \{a_{x'}\}$ . Take  $A_r = \{a_{x'}\}$ .  $\square$

**Lemma 3.10.**  $\varphi : L \rightarrow M$  is  $\alpha$ -complete if and only if  $Y(\varphi) : Y(M) \rightarrow Y(L)$  is an  $\alpha$ -SpFi morphism.

**Proof.** Let  $\varphi$  be  $\alpha$ -complete. For  $G \in \mathcal{G}_\alpha(Y(L))$  take an  $A_1 \subset L^+$  as in Lemma 3.9(c). Since  $\varphi$  is  $\alpha$ -complete, and  $\varphi(a) = a \circ Y(\varphi)$  for all  $a \in A$  (Theorem 3.3(a)), it follows that  $1 = \varphi(1) = \bigvee^M \{\varphi(a) : a \in A\} = \bigvee^M \{a \circ Y(\varphi) : a \in A\}$ . Applying Lemma 3.9(a) we know that  $P_1 = \{x \in Y(M) : 1 = \bigvee_{A_1}^{\mathbb{R}} a \circ Y(\varphi)(x)\}$  is dense in  $Y(M)$ . Now, because

$x \in P_1$  implies that  $x \in Y(\varphi)^{-1}(\text{coz}(a))$  for some  $a \in A$ , we have  $P_1 \subset Y(\varphi)^{-1}(\bigcup_A \text{coz}(a)) = Y(\varphi)^{-1}(G)$ . So  $Y(\varphi)^{-1}(G)$  is dense in  $Y(M)$ . Therefore  $Y(\varphi) \in \alpha\text{-SpFi}$  (Proposition 3.8(b)).

On the other hand, suppose  $Y(\varphi)$  is an  $\alpha\text{-SpFi}$  morphism and let  $\mathbf{1} = \bigvee^L A$  with  $|A| < \alpha$ . Applying Lemma 3.9 we know that  $P_1 = \{x \in Y(L) : 1 = \bigvee_A a(x)\} \in \mathcal{G}_\alpha(Y(L))_\delta$ . For  $x \in Y(M)$ , if  $Y(\varphi)(x) \in P_1$ , then  $1 = \bigvee_A a \circ Y(\varphi)(x) = \bigvee_A \varphi(a)(x)$ . Hence  $Y(\varphi)^{-1}(P_1) \subset \{y \in Y(M) : 1 = \bigvee_A \varphi(a)(y)\}$  is dense (Proposition 3.8(c)). So  $\varphi$  is  $\alpha$ -complete (Proposition 2.1(b)).  $\square$

**Lemma 3.11.**  $\varphi : L \rightarrow M$  is epic in  $\mathcal{W}(\alpha)$  if and only if  $Y(\varphi) : Y(M) \rightarrow Y(L)$  is monic in  $\alpha\text{-SpFi}$ .

**Proof.** Let  $\varphi$  be epic in  $\mathcal{W}(\alpha)$  and suppose for  $i = 1, 2$  that  $f_i : X \rightarrow Y(M)$  are  $\alpha\text{-SpFi}$  morphisms, and  $Y(\varphi) \circ f_1 = Y(\varphi) \circ f_2$ .

We utilize a result of Gleason:

For each  $X$ , there is an essentially unique pair  $(EX, \pi)$  such that  $\pi : EX \twoheadrightarrow X$  is  $\infty$ -irreducible (Definition 3.14) and  $EX$  is extremally disconnected [12].  $(EX, \pi)$  is called the *absolute* of  $X$  [25]. The map  $\pi$  is also an  $\infty\text{-SpFi}$  morphism (Lemma 3.15). Since  $EX$  is extremally disconnected,  $D(EX)$  is a  $\mathcal{W}$  object [17]. For  $i = 1, 2$ , let  $h_i = f_i \circ \pi$ . The  $h_i$  are  $\alpha\text{-SpFi}$  morphisms. Define  $\gamma_i : M \rightarrow D(EX)$ , for  $i = 1, 2$ , by  $\gamma_i(b) = b \circ h_i$  for  $b \in M$ . This definition makes sense because  $M \subseteq D(Y(M))$  and so  $b^{-1}(\mathbb{R}) \in \mathcal{G}_{\omega_1}(Y(M))$  (i.e.,  $\gamma_i(b) \in D(X)$  because  $\gamma_i(b)^{-1}(\mathbb{R}) = h_i^{-1}(b^{-1}(\mathbb{R}))$  is dense in  $EX$  by virtue of the map  $h_i$  being an  $\alpha\text{-SpFi}$  morphism). Clearly,  $\gamma_i$  is a homomorphism, and by Theorem 3.3(a),  $Y(\gamma_i) = h_i$ . Therefore the  $\gamma_i$  are  $\alpha$ -complete (Lemma 3.10). Moreover,  $Y(\gamma_1 \circ \varphi) = Y(\varphi) \circ Y(\gamma_1) = Y(\varphi) \circ Y(\gamma_2) = Y(\gamma_2 \circ \varphi)$ . Therefore because  $Y$  is faithful (Theorem 3.3(b)),  $\gamma_1 \circ \varphi = \gamma_2 \circ \varphi$ . Hence  $\gamma_1 = \gamma_2$ , and  $h_1 = Y(\gamma_1) = Y(\gamma_2) = h_2$ .  $f_1 = f_2$  because  $\pi$  is onto, so  $Y(\varphi)$  is monic in  $\alpha\text{-SpFi}$ .

Now suppose  $Y(\varphi)$  is monic in  $\alpha\text{-SpFi}$  and suppose for  $i = 1, 2$ , that  $\gamma_i : M \rightarrow N$  are  $\alpha$ -complete with  $\gamma_1 \circ \varphi = \gamma_2 \circ \varphi$ . Then  $Y(\varphi) \circ Y(\gamma_1) = Y(\varphi) \circ Y(\gamma_2)$ . Since  $Y(\gamma_i)$  is an  $\alpha\text{-SpFi}$  morphism (Lemma 3.10), we have by our assumption that  $Y(\gamma_1) = Y(\gamma_2)$ . And so  $\gamma_1 = \gamma_2$ . Therefore  $\varphi$  is epic.  $\square$

**Theorem 3.12** [5]. *The following are equivalent:*

- (a)  $X$  is  $\alpha$ -quasi- $F$  (Definition 1.5).
- (b) Each  $G \in \mathcal{G}_\alpha(X)_\delta$  is  $C^*$ -embedded in  $X$ .
- (c)  $C(X)$  is  $\alpha$ -Dedekind complete.

**Theorem 3.13.** *If  $L$  is  $\alpha$ -Dedekind complete, then  $Y(L)$  is  $\alpha$ -quasi- $F$ .*

**Proof.** Let  $K \subset Y(L)$  be a dense  $\alpha$ -Lindelöf subspace and let  $h \in C^*(K)$ . Without loss of generality, assume  $0 \leq h \leq 1$ . For each  $n \in \mathbb{N}$  and  $p \in K$  there exists a neighborhood,  $V_p^n$ , of  $p$  in  $Y(L)$  such that  $\max h[\bar{V}_p^n \cap K] - \min h[\bar{V}_p^n \cap K] < n^{-1}$ . Choose a neighborhood,  $U_p^n$ , of  $p$  in  $Y(L)$  so that  $\bar{U}_p^n \subset V_p^n \subset \bar{V}_p^n$ . Let  $M = \max h[\bar{V}_p^n \cap K]$

and  $m = \min h[\bar{V}_p^n \cap K]$ . Since  $L^* \subseteq C(Y(L))$  separates the points of  $Y(L)$  there are functions  $a_p^n, b_p^n \in L^*$  such that:

- (i)  $0 \leq a_p \leq m; M \leq b_p \leq 1$ .
- (ii)  $a_p^n[\bar{U}_p^n] = m; a_p^n[Y(L) \setminus V_p^n] = 0$ .
- (iii)  $b_p^n[\bar{U}_p^n] = M; b_p^n[Y(L) \setminus V_p^n] = 1$ .

It follows that  $a_{p'}^n(y) < h(y) < b_p^n(y)$  for all  $n, n' \in \mathbb{N}$  and  $p, p', y \in K$ .

Because  $K$  is  $\alpha$ -Lindelöf we have, for each  $n \in \mathbb{N}$ , a subset  $P' \subset K$  such that  $|P'| < \alpha$  and  $K = \bigcup_{p' \in P'} \{U_{p'}^n\}$ . Let  $A = \{a_{p'}^n: n \in \mathbb{N}, p' \in P'\}$  and  $B = \{b_{p'}^n: n \in \mathbb{N}, p' \in P'\}$ . It follows that  $|A|, |B| < \alpha$  and  $B \geq A$ . We claim  $0 = \bigwedge^L (B - A)$ . For  $K \subseteq \{x \in X: 0 = \bigwedge_{A,B}^{\mathbb{R}} (b(x) - a(x))\}$ , and since  $K$  is dense in  $Y(L)$  we have, as in Lemma 3.9(a), that  $0 = \bigwedge^L (B - A)$ . Therefore, because  $L$  is  $\alpha$ -Dedekind complete, there is a  $g \in L^* \subseteq C(Y(L))$  with  $g = \bigwedge B = \bigvee A$ . Clearly,  $g|_K = h$ , hence  $K$  is  $C^*$ -embedded in  $Y(L)$ .  $\square$

Definition 3.14 below shows how  $Y$  links the notions of  $\alpha$ -density and  $\alpha$ -irreducibility. Lemma 3.15 is a slight generalization of the equivalence of 3.12(a) and 3.12(g) in [5].

**Definition 3.14.** Let  $f: X \rightarrow Y$ .  $f$  is said to be  $\alpha$ -irreducible if for each  $U \in \text{Coz}(X)$ , there is a  $V \in \text{Coz}_\alpha(Y)$  such that  $f^{-1}(V) \subset U$  and  $f^{-1}(V) = \bar{U}$ .

**Lemma 3.15.** An embedding  $\varphi: L \hookrightarrow M$  is  $\alpha$ -dense if and only if  $Y(\varphi): Y(M) \rightarrow Y(L)$  is  $\alpha$ -irreducible.

**Proof.** Suppose  $Y(\varphi)$  is  $\alpha$ -irreducible,  $b \in M^+$ , and  $U = \text{coz}(b) \cap b^{-1}(\mathbb{R})$ . Since  $U$  is  $\omega_1$ -Lindelöf ( $b^{-1}(\mathbb{R}) \in \text{Coz}(Y(M))$ ) and  $b$  does not take the value  $\pm\infty$  anywhere on  $U$ , there is for each  $n \in \mathbb{N}$  a family  $\{U_k^n: k \in \mathbb{N}\} \subset \text{Coz}(Y(M))$  such that  $U = \bigcup_k U_k^n = \bigcup_k \bar{U}_k^n$ , and  $\max b[\bar{U}_k^n] - \min b[\bar{U}_k^n] \leq n^{-1}$  for each  $k \in \mathbb{N}$ .

The indices will now get a bit hairy.

Since  $Y(\varphi)$  is  $\alpha$ -irreducible, we have for each  $(n, k) \in \mathbb{N} \times \mathbb{N}$ , a  $V_k^n \in \text{Coz}_\alpha(Y(L))$  such that  $Y(\varphi)^{-1}(V_k^n)$  is dense in  $U_k^n$ . Also, since for each  $(n, k)$ ,  $V_k^n$  is  $\alpha$ -Lindelöf, there is a family,  $\{V_k^n(i): i \in I, |I| < \alpha\} \subset \text{Coz}(Y(L))$ , such that  $V_k^n = \bigcup_i \bar{V}_k^n(i)$ . Finally, there is a family of functions in  $L$ ,  $\{a_k^n(i): (i, n, k) \in I \times \mathbb{N} \times \mathbb{N}\}$ , such that for each  $(i, n, k)$ :

- (i)  $0 \leq a_k^n(i) \leq \min b[\bar{U}_k^n]$ .
- (ii)  $a_k^n(i)[\bar{V}_k^n(i)] = \min b[\bar{U}_k^n]$ .
- (iii)  $a_k^n(i)[Y(L) \setminus V_k^n] = 0$ .

Upon careful reflection (and recalling that  $a_k^n(i) \circ Y(\varphi)(y) = \varphi(a_k^n(i))(y)$ ) one will see that

$$\bigcap_n \left( \bigcup_k \{Y(\varphi)^{-1}(V_k^n) \cup (Y(M) \setminus U)^0\} \right) \subset \left\{ y \in Y(M): \bigvee_{i,n,k} a_k^n(i) \circ Y(\varphi)(y) = b(y) \right\}.$$

Invoking the Baire category theorem, one can see that the former, and hence, the latter set is dense in  $Y(M)$  because for each  $n \in N$ ,  $\bigcup_k Y(\varphi)^{-1}(V_k^n)$  is dense in  $U$ ; so  $\bigcup_k \{Y(\varphi)^{-1}(V_k^n) \cup (Y(M) \setminus U)^o\}$  is dense and open in  $Y(M)$ . Therefore

$$\begin{aligned} & \left\{ y \in Y(M) : \bigvee_{i,n,k} a_k^n(i) \circ Y(\varphi)(y) = b(y) \right\} \\ &= \left\{ y \in Y(M) : \bigvee_{i,n,k} \varphi(a_k^n(i))(y) = b(y) \right\} \end{aligned}$$

is dense in  $Y(M)$ . Applying Lemma 3.9(a), we have that  $b = \bigvee_{i,n,k}^L \varphi(a_k^n(i))$ . Since  $|I \times \mathbb{N} \times \mathbb{N}| < \alpha$ , it follows that  $\varphi$  is  $\alpha$ -dense.

On the other hand, suppose  $\varphi$  is an  $\alpha$ -dense embedding. Let  $U \in \text{Coz}(Y(M))$ . We can assume without loss of generality that  $U = \text{coz}(b)$  for some  $b \in M^+$ . Since  $\varphi$  is  $\alpha$ -dense, there is a  $A \subset L$  with  $|A| < \alpha$  such that  $b = \bigvee^M \varphi[A]$ . If we let  $P_b = \{y \in Y(M) : \bigvee_A^M \varphi(a)(y) = b(y)\}$ , then  $P_b$  is dense in  $Y(M)$  (Lemma 3.9(a)). We claim that for  $V = (\bigcup_A \text{coz}(a)) \in \text{Coz}_\alpha(Y(L))$ , we have that  $Y(\varphi)^{-1}(V)$  is dense in  $U$ . Now

$$\begin{aligned} \text{coz}(b) \cap P_b &\subset \bigcup_A \text{coz}(\varphi(a)) \\ &= \bigcup_A \text{coz}(a \circ Y(\varphi)) = Y(\varphi)^{-1}(V) \subset \text{coz}(b) = U. \end{aligned}$$

Since  $P_b$  is dense in  $Y(M)$ ,  $Y(\varphi)^{-1}(V)$  is dense in  $U$ .  $\square$

The following is a generalization to arbitrary  $\alpha$  of a lemma of Weinberg [29].

**Corollary 3.16.**  *$f: X \rightarrow Y$  is  $\alpha$ -irreducible if and only if  $f': C(Y) \rightarrow C(X)$  is  $\alpha$ -dense. Moreover, if  $f$  is  $\alpha$ -irreducible, then  $f'$  is an  $\alpha$ -SpFi morphism.*

**Proof.**  $Y(f') = f$ . Apply Definition 3.14, Corollary 2.7, and Lemma 3.10.  $\square$

Note, Lemma 3.15 follows directly from [5, 3.12].

#### 4. The $\alpha$ -quasi- $F$ cover

In this section all spaces are assumed to be compact and Hausdorff. Theorem 4.1 below is a synthesis of results taken from [5, 9, 18, 24]. See also [13, 22, 23, 27]. Recall the definition of the minimum  $P$  cover of  $X$  (Definition 1.6). Here we take  $P = \alpha$ -quasi- $F$ .

**Theorem 4.1.** *For each  $X$ , there exists a minimum  $\alpha$ -quasi- $F$  cover, denoted by  $(QF_\alpha X, q_\alpha)$ . Moreover,  $q_\alpha$  is  $\alpha$ -irreducible; and any  $\alpha$ -quasi- $F$  cover  $(Y, f)$  of  $X$  with  $f$   $\alpha$ -irreducible is essentially unique.*

**Theorem 4.2.** *Let  $(i, C(X)_\alpha^\wedge)$  be the  $\alpha$ -Dedekind complete epireflection of  $C(X)$  in  $\mathcal{W}(\alpha)$  (Theorem 2.19). Then  $(Y(C(X)_\alpha^\wedge), Y(i))$  is essentially identical to the  $\alpha$ -quasi- $F$  cover of  $X$ .*



**Proof.** Apply Theorem 3.13, Definition 3.14, and Theorem 4.1.  $\square$

We now give the main result of this section. We say  $(Y, f)$  is an  $\alpha$ -irreducible preimage of  $X$  if  $f: X \rightarrow Y$  is  $\alpha$ -irreducible. Recall that in an abstract category  $\mathcal{B}$ , a full subcategory  $\mathcal{A}$  is called *monocoreflective* if for each  $B \in |\mathcal{B}|$  there is an  $A_B \in |\mathcal{A}|$  and a monic  $m_B: A_B \rightarrow B$  such that whenever  $f: A \rightarrow B$  and  $A \in |\mathcal{A}|$  there is an  $\bar{f}: A \rightarrow A_B$  with  $f = m \circ \bar{f}$ .

**Theorem 4.3.** (a) Suppose  $(Y, f)$  is an  $\alpha$ -irreducible preimage of  $X$ ;  $h: Z \rightarrow X$  is  $\alpha$ -SpFi; and  $Z$  is  $\alpha$ -quasi- $F$ . Then there exists a unique  $\alpha$ -SpFi morphism  $\bar{h}: Z \rightarrow Y$  with  $h = f \circ \bar{h}$ .

(b) In  $\alpha$ -SpFi, the full subcategory of  $\alpha$ -quasi- $F$  spaces is monocoreflective, and  $(QF_\alpha X, q_\alpha)$  is the  $\alpha$ -quasi- $F$  monocoreflection of  $X$ .

(c)  $(QF_\alpha X, q_\alpha)$  is the maximal  $\alpha$ -irreducible preimage of  $X$ . That is, if  $(Y, f)$  is an  $\alpha$ -irreducible preimage of  $X$ , then there is an  $\hat{f}: QF_\alpha X \rightarrow Y$  such that  $q_\alpha = f \circ \hat{f}$ . It then follows that  $\hat{f}$  must be  $\alpha$ -irreducible and  $QF_\alpha X \cong QF_\alpha Y$ .

**Proof.** (a) Let  $(Y, f)$  be as above. By Lemma 3.15,  $(f', C(Y))$  is an  $\alpha$ -dense extension of  $C(X)$ . By Lemma 3.10 and Theorem 3.12 respectively, if  $h: Z \rightarrow X$  is an  $\alpha$ -SpFi morphism and  $Z$  is  $\alpha$ -quasi- $F$ , then  $h': C(X) \rightarrow C(Z)$  is  $\alpha$ -complete and  $C(Z)$  is  $\alpha$ -Dedekind complete. Hence by Theorem 2.16, there is a unique  $\alpha$ -complete  $\varphi: C(Y) \rightarrow C(Z)$  such that  $f' = \varphi \circ h'$ . Applying the Yosida functor we get that  $Y(\varphi): Z = Y(C(Z)) \rightarrow Y(C(Y)) = Y$  is an  $\alpha$ -SpFi morphism and  $h = Y(h') = Y(f') \circ Y(\varphi) = f \circ Y(\varphi)$ . Take  $\bar{h} = Y(\varphi)$ .

(b) It follows from Proposition 2.8, Lemma 3.11 and Definition 3.14 that  $q_\alpha$  is monic in  $\alpha$ -SpFi. Apply (a) here and Theorem 4.1.

(c) By (a) here we have a map  $\hat{f}: QF_\alpha X \rightarrow Y$  such that  $q_\alpha = \hat{f} \circ f$ . It is not hard to see that since  $q_\alpha$  and  $f$  are  $\alpha$ -irreducible, then  $\hat{f}$  must also be  $\alpha$ -irreducible. Finally, to show  $QF_\alpha X \cong QF_\alpha Y$ , apply Theorem 4.1.  $\square$

Theorem 4.3(b) was proved independently by A. Molitor in [23], and Theorem 4.3(c) has been demonstrated in [18] when  $\alpha = \omega_1$ .

## Acknowledgement

The author is indebted to A.W. Hager and the referee for many helpful comments and suggestions. The main body of this paper was taken from the authors thesis, written at Wesleyan University.

## References

- [1] R.N. Ball, Convergence and Cauchy structures on lattice-ordered groups, Trans. Amer. Math. Soc. 259 (1980) 357–392.

- [2] R.N. Ball and A.W. Hager, Characterization of epimorphisms in archimedean  $l$ -groups and vector lattices, in: A. Glass and W.C. Holland, eds., *Lattice-Ordered Groups, Advances and Techniques* (Kluwer, Dordrecht, 1989).
- [3] R.N. Ball and A.W. Hager, Epicompletion or archimedean  $l$ -groups and vector lattices with weak unit, *J. Austral. Math. Soc. Ser. A* 48 (1990) 25–56.
- [4] R.N. Ball, A.W. Hager and A.J. Macula, An  $\alpha$ -disconnected space has no proper monic preimage, *Topology Appl.* 37 (1990) 141–151.
- [5] R.N. Ball, A.W. Hager and C. Neville, The  $\kappa$ -ideal completion of an archimedean  $l$ -group and the  $\kappa$ -quasi- $F$  cover of a compact space, in: R.M. Shortt, ed., *Proceedings Northeast Topology Conference*, Wesleyan University, 1988, *General Topology and its Applications* 123 (Dekker, New York, 1990) 7–50.
- [6] A. Bigard, K. Keimel and S. Wolfenstein, *Groupes et Anneaux Reticules*, *Lecture Notes in Mathematics* 608 (Springer, Berlin, 1977).
- [7] G. Birkhoff, *Lattice Theory*, *American Mathematical Society Colloquium Publications* (Amer. Math. Soc., Providence, RI, 3rd ed., 1973).
- [8] W.W. Comfort and S. Negropontis, *Continuous Pseudometrics*, *Lecture Notes in Pure and Applied Mathematics* 14 (Dekker, New York, 1975).
- [9] F. Dasheill, A.W. Hager and M. Henriksen, Order-Cauchy completions of rings and vector lattices of continuous functions, *Canad. J. Math.* 32 (1980) 657–685.
- [10] E. DeJonge and A.C.M. Van Rooij, *Introduction to Riesz Spaces*, *Mathematical Centre Tracts* 78 (Math. Centre, Amsterdam, 1977).
- [11] L. Gillman and M. Jerison, *Rings of Continuous Functions* (Van Nostrand Reinhold, Princeton, NJ, 1960).
- [12] A.M. Gleason, Projective topological spaces, *Illinois J. Math.* 2 (1958) 482–489.
- [13] A.W. Hager, Minimal covers of topological spaces, in: *Proceedings Northeast Topology Conference*, 1987, *Annals of the New York Academy of Sciences* 552 (New York Acad. Sci., New York, 1989) 44–59.
- [14] A.W. Hager, Boolean algebras with  $\alpha$ -complete morphisms, *Manuscript*.
- [15] A.W. Hager and L.C. Robertson, Representing and ringifying a Riesz space, *Symposia Mathematica* 21 (Academic Press, New York, 1977) 411–431.
- [16] A.W. Hager and L.C. Robertson, On embedding into a ring of an archimedean lattice-ordered group, *Canad. J. Math.* 31 (1979) 1–8.
- [17] M. Henriksen and D.G. Johnson, On the structure of a class of archimedean lattice-ordered algebras, *Fund. Math.* 50 (1961) 73–94.
- [18] M. Henriksen, J. Vermeer and R.G. Woods, Quasi- $F$  covers of Tychonoff spaces, *Trans. Amer. Math. Soc.* 303 (1987) 779–803.
- [19] H. Herrlich and G. Strecker, *Category Theory* (Allyn and Bacon, Boston, MA, 1973).
- [20] A.J. Macula, Thesis, Wesleyan University (1989).
- [21] A.J. Macula, Free  $\alpha$ -extensions of an archimedean vector lattice and their topological duals, *Trans. Amer. Math. Soc.*, to appear.
- [22] J. Madden,  $\kappa$ -frames, *Manuscript*.
- [23] A. Molitor, Quasi- $F$ -like covers of a compact Hausdorff space, in: R.M. Shortt, ed., *Proceedings Northeast Topology Conference*, Wesleyan University, 1988, *General Topology and its Applications* 123 (Dekker, New York, 1990) 219–226.
- [24] C.W. Neville, Quasi- $F_\eta$  spaces, *Manuscript*.
- [25] J.R. Porter and R.G. Woods, *Extensions and Absolutes of Hausdorff Spaces* (Springer, New York, 1988).
- [26] R. Sikorski, *Boolean Algebra* (Springer, Berlin, 3rd ed., 1969).
- [27] J. Vermeer, On irreducible preimages, *Topology Proc.* 9 (1984) 173–189.
- [28] R.C. Walker, *The Stone-Čech Compactification* (Springer, Berlin, 1974).
- [29] E. Weinberg, Thesis, *Perdue University* (1961).
- [30] R.G. Woods, Covering properties and coreflective subcategories, in: *Proceedings CCNY Conference on Limits 1987*, *Annals of the New York Academy Science* (New York Acad. Sci., New York, to appear).
- [31] K. Yosida, On the representation of a vector lattice, *Proc. Imp. Acad. Tokyo* 18 (1942) 339–342.
- [32] W.A.J. Luxemburg and A.C. Zaanen, *Riesz Spaces I* (North-Holland, Amsterdam, 1971).